Obsidian® ASG Application Kit

Quick Reference Card




* APL 400 Series

Local language versions are available for download on www.vivostat.com/grcs

PART 1 – PREPARE DISPOSABLE APPLICATOR DEVICE AND APPLICATOR UNIT


Turn ON Applicator Unit on the back and connect foot switch




1. Place 21g needle just inside pH 10 vial, draw fluid into 1 ml syringe



2. Transfer disposable applicator + pH-paper out of sterile to sterile area



3. Transfer syringe holder area




4. Insert pH 10 and Obsidian syringes into syringe holder



5. Rotate syringe holder back and forth (x10)



6. When indicated in display, open shield



7. Insert and position syringe holder



8. Press syringe 9. Close shield holder into cavity using two fingers



CAUTION:

- 1) To avoid air bubbles impacting mixture negatively do NOT push needle to the end of pH 10 vial in Step 1
- 2) Follow Step 5 carefully for platelets not to settle at the bottom, as they are essential in the healing process
- 3) If +30 min. pass after step 5 rotate syringe holder again: Press button on top of Applicator Unit for +4 sec., take out

PART 2 (OPTIONAL) - LOAD CATHETER IN STERILISED ENDOSCOPIC APPLICATOR HANDLE



1. Place endoscopic catheter tip at the end of handle



endoscopic catheter entirely through



2. Load and push 3. Adjust endoscopic catheter for the right position - all angles are possible



4. To lock position push catheter up until it "clicks"

PART 3 – LOAD SYRINGES, PRIME AND INITIATE APPLICATION



1. Press button to start loading syringes



2. Press foot switch once to prime




3. Point to lower corner of pH-paper while auto-priming



4. Autopriming stops automatically and ml + spray mode is displayed



5. Verify correct mixture on pHpaper (from liquid to viscous in 10-30 sec.)



6. Ready for application, follow separate Obsidian® ASG Application Instruction