

Experienced use across Europe

Good polymerization

- Amikacin
- Augmentin
- Bacitracin (Baneocin)
- Biklin
- Cefrom
- Ciprofloxacin
- Ciprofloxacin + Colistin
- Colistin (Promixin)
- Flucloxacillin
- Gentamycin
- Penem – certain kinds
- Tazobactam (Piperacillin/Tazonam)
- Teicoplanin
- Tobramycin (Nebcina)
- Tobramycin + Colistin
- Unasyn (Ampicillin/Sulbactam)
- Vancomycin

No polymerization

- Cefepim
- Clindamycin/Dalacin
- Cubicin
- Ertapenem

Experienced use – Medical University of Vienna¹

Good polymerization

- Amikacin
- Teicoplanin
- Teicoplanin + amikacin

Dosage/ml aqua

250 mg/ml
400 mg/1.5 ml
400 mg + 2.6 ml

Medication reconstitution in water

prepared solution
5 min.
10 min.

Slow polymerization

- Ampicillin/Sulbactam

Dosage/ml aqua

1.5 g/3 ml

Medication reconstitution in water

5 min.

No polymerization

- Ertapenem

Dosage/ml aqua

1 g/1.5 ml

Medication reconstitution in water

5 min.

1: Dr. M. Vossen, Medical University of Vienna